

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 808 896 A2

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

26.11.1997 Bulletin 1997/48

(51) Int. Cl.<sup>6</sup>: C12N 5/00

(21) Application number: 97108184.9

(22) Date of filing: 21.05.1997

(84) Designated Contracting States:  
DE FR GB

(30) Priority: 21.05.1996 JP 125533/96

(71) Applicant:  
The Institute of Physical and Chemical Research  
(RIKEN)  
Wako-shi, Saltama 351-01 (JP)

(72) Inventors:

- Kusakabe, Moriaki  
Koyadai, 3-1-1, Tsukuba-shi, Ibaragi 305 (JP)
- Kamon, Toshio  
20-14, Aobadai 2-chome, Meguro-ku Tokyo 153 (JP)

(74) Representative:

Godemeyer, Thomas, Dr.  
Hauptstrasse 58  
51491 Overath (DE)

(54) **Embryonic stem cells from mice**

(57) Embryonic stem cells derived from an inbred mouse strain selected from the group consisting of C3H/HeN, C57BL/6N, DBA/1J, and BALB/c strains, for example, C-2 cell having the accession number FERM BP-5933 and C-6 cell having the accession number FERM BP-5934 are disclosed. The embryonic stem cells of the present invention are derived from genetically complete inbred strains, and therefore, they are extremely useful for close genetic researches. For example, they are useful for the production of knockout mice as well as for analyses of cell genealogical tree and mechanism of embryonic stem cell establishment.

EP 0 808 896 A2

**BEST AVAILABLE COPY**

## Description

## BACKGROUND OF THE INVENTION

## Field of the Invention

The present invention relates to novel embryonic stem cells derived from mice.

## Related Art

Embryonic stem cells, sometimes referred to as ES cells, are derived from inner cell mass (ICM) of fertilized eggs in blastocyst phase, and can be cultured and maintained *in vitro* while being kept in undifferentiated state. Embryonic stem cells are extremely useful biological materials for preparing transgenic animals. For example, a gene knockout mouse in which a specific gene is inactivated can be produced by replacing an active gene in an embryonic stem cell chromosome with an inactivated gene by means of a homologous recombination system.

Embryonic stem cells derived from mice, hamsters, and pigs were previously reported. However, processes for establishing embryonic stem cells have not been sufficiently developed, and the sorts of embryonic stem cells are undeniably limited compared to the numbers of mouse strains that have been developed for wide variety of numerous purposes. Currently, researchers most widely use the cells derived from mouse strain 129/Sv as embryonic stem cells. However, the mice of 129/Sv are not always sufficiently bred, and accordingly, if a gene knockout mouse is established, the mouse must be disadvantageously made into a hybrid by crossbreeding with other strains. There is also a problem that the origin of embryonic stem cells established from the strain 129/Sv cannot be completely specified from a genetic viewpoint, since various kinds of substrains are involved in the strain.

Function of a certain gene is generally influenced by other genetic backgrounds in a whole-body level. Accordingly, for close comparative researches by means of gene knockout mice, it is indeed ideal to establish embryonic stem cells from an inbred mouse strain, whose basic data has sufficiently been accumulated, and compare the influence of the knockout of a embryonic stem cells from an inbred mouse strain, whose basic data has sufficiently been accumulated, and compare the influence of the knockout of a gene of interest with results obtained from original strain. For these reasons, it has been desired to establish novel embryonic stem cells from inbred mouse strains. However, this class of embryonic stem cells has not been reported yet. As embryonic stem cells derived from mouse strains, embryonic undifferentiated cells disclosed in Japanese Patent Unexamined Publication (KOKAI) No. (Hei) 5-328878/1993 and the like are known. However, these cells are characterized as embryonic stem cells derived from F1 hybrid, and are not those established from inbred strains.

Accordingly, an object of the present invention is to provide novel embryonic stem cells. More specifically, the object of the present invention is to provide embryonic stem cell established from inbred mouse strains.

## DESCRIPTION OF THE INVENTION

The inventors of the present invention conducted various researches to achieve the foregoing object, and as a result, they succeeded in establishing novel embryonic stem cells from inbred mouse strains such as C3H/HeN.

The present invention thus provide embryonic stem cells derived from mouse strains C3H/HeN, C57BL/6N, DBA/1J, and BALB/c.

## DETAILED EXPLANATION OF THE INVENTION

## Explanation of the Preferred Embodiments

The embryonic stem cell of the present invention were established from mouse inbred strains of C3H/HeN, C57BL/6N, DBA/1J, and BALB/c. Among these inbred strains, C3H/HeN, C57BL/6N and DBA/1J are genetically complete inbred strains. As for BALB/c strain, some substrains, e.g., BALB/cA and BALB/cC are known, however, the substrains are genetically very close to each other. These mouse strains are widely and conventionally used and can easily be obtained.

For example, the embryonic stem cells of the present invention can be established as follows: Male and female mice of the above strains are allowed to natural mating, and on the next day, the female mice having a vaginal plug are assigned as day zero (0) of pregnancy. Blastocysts are removed from uteri of the female mice of day three (3) of pregnancy and then cultured. After hatching, the cells are transferred on feeder cells and cultivation is continued for 2 or 3 days. The feeder cells can be prepared, for example, by treating fibroblasts of 14th-day embryos of mouse BALB/cA strain with Mitomycin C. An inner cell mass (ICM) become rising is exfoliated and made into single cell masses in approximately 0.25% trypsin droplets. The small cell masses are transferred on the feeder cells in a 24-well plate as the

primary culture. The successive treatments include the following steps: The culture medium is changed every day from the next day, and each colonies are separately picked up after 2 to 5 days and made into single cell masses with trypsin treatments. These dissociated cells are then placed again on the feeder cells. This procedure is repeated up to tenth subculture to allow the establishment of embryonic stem cells.

## Examples

The present invention will be explained more specifically by referring to the examples. However, the scope of the present invention is not limited to these examples.

### (1) Collection of ovum

Male and female mice of respective C3H/HeN, C57BL/6N, or DBA/1J strains were allowed to natural breed, and uteri of the mice on the 3rd pregnant day were removed and washed with M2 buffer. As for BALB/c, F1 embryos between substrains that were formed by crossbreeding of BALB/cA and BALB/cCr strains were used. Embryos were washed out from the uteri with M2 buffer using a syringe equipped with a perfusion needle (30 gage). The blastocysts with zona pellucida were washed twice with M2, and then cultured in ESM medium in an incubator under 5% CO<sub>2</sub> at 37°C for 30 hours until the embryos hatched [a droplet of the ESM medium having a diameter of 5 mm was formed and overlaid with mineral oil (Sigma), and then preliminarily incubated in an incubator under 5% CO<sub>2</sub> at 37°C]. The hatched blastocysts were transferred on feeder cells, which had been obtained from 14th-day embryos of BALB/cA mouse and treated with Mitomycin C (10 µg/ml in DMEM), and then cultured in ESM medium. Risings of ICM were observed in the hatched embryos after two or three days.

### (2) Isolation and unbinding of ICM

The rising ICMs were picked up by suction using a 50 µl micropipette containing a small volume of PBS(-) sucked beforehand and twice washed with PBS(-), and then transferred into a droplet of a 0.25% trypsin solution and incubated at room temperature for 1 to 1.5 minutes. Before the cell mass was unbound, a small volume of ESM was added into the trypsin droplet using a micropipette to inactivate trypsin. The cell mass was divided into appropriately small pieces in separate EMS droplets, and each of the masses was transferred on feeder cells treated with Mitomycin C (10 µg/ml in DMEM). The resulted mass was assigned to the primary culture, and the culture medium was changed every day from the next day. For the second and later subcultures, cell masses were completely unbound to obtain individual cells in ESM after the treatment of trypsin for 40 to 60 seconds, and the cells were transferred on the feeder cells in a similar manner.

(a) ESM culture medium: 400 ml of DMEM (0.012 g of potassium penicillin G; 0.02 g of streptomycin sulfate; 0.6 g of sodium bicarbonate; and 4 g of DMEM were dissolved in water in a final volume of 400 ml); 2 ml of NEAA (non-essential amino acid, × 100); 4 ml of nucleoside stock solution (3 mM adenosine; 3 mM guanosine; 3 mM uridine; 3 mM cytidine; and 1 mM thymidine); 0.4 ml of 2-mercaptoethanol stock solution (5 ml of DMEM and 35 µl of 2-mercaptoethanol); 50 µl of LIF (leukemia inhibiting factor); 1.85 g of glucose; and 100 ml of FCS were mixed.

(b) 0.25% trypsin solution; trypsin (0.25 g); EDTA.2Na (0.037 g); and 1 × phosphate buffered saline (-) (100 ml).

(c) M2 buffer: 15.0 ml of stock solution A (11.068 g of NaCl; 0.712g of KCl; 0.324 g of KH<sub>2</sub>PO<sub>4</sub>; 0.586g of MgSO<sub>4</sub>·7H<sub>2</sub>O; 8.698 ml of sodium lactate (60% syrup); 2.000 g of glucose; 0.120 g of potassium penicillin G; and 0.100 g of streptomycin sulfate in 200 ml); 2.4 ml of stock solution B (1.051 g of NaHCO<sub>3</sub> and 0.005 g of phenol red in 50 ml); 1.5 ml of stock solution C (0.036 g of sodium pyruvate in 10 ml); 1.5 ml of stock solution D (1.260 g of CaCl<sub>2</sub>·2H<sub>2</sub>O in 50ml); 12.6 ml of stock solution E (5.958 g of HEPES and 0.010 g of phenol red in 100ml; adjusted to pH 7.4 with NaOH); and 600 mg of BSA were mixed and adjusted to 150 ml with distilled water.

### (3) Selection of embryonic stem cells

After 2 to 5-day cultivation, cell populations of various shapes were observed. Among them, flat colonies formed by morphologically uniform small cells were picked up as embryonic stem cell populations, and the colonies were treated with trypsin for 40 to 60 seconds and then subcultured on feeder cells freshly prepared as described in the above section (2). Subcultures were repeated up to tenth subculture to establish embryonic stem cells of the present invention. The cells were deposited on May 17, 1996 at the National Institute of Bioscience and Human-Technology Agency of Industrial Science and Technology (1-3, Higasi-1-chome, Tsukuba-shi, Ibaraki-ken 305 Japan) and the original depositions were transferred to the International Depositary Authority under Budapest treaty identified above on May 1, 1997.

Table 1

Embryonic stem cell	Mouse strain	Accession number
C-6	C3H/HeN	FERM BP-5934
C-2	C3H/HeN	FERM BP-5933
Cr/A-3	BALB/c	FERM BP-5932
DB1	DBA/1J	FERM BP-5931
B6-26	C57BL/6N	FERM BP-5935

All of these embryonic stem cells were positive in alkali phosphatase activity staining. In addition, among these embryonic stem cells, C-6, C-2, Cr/A-3, and DB1 were evaluated as for the ability of differentiation into germ cells by producing chimera mice according to the injection method or the aggregation method. The results are shown below.

Table 2

ES cell-line <sup>1)</sup>	Chimera mouse	Recipient embryo strain	Mating strain	Number % litter	Originate from stem cells <sup>2)</sup>	Originate from recipient embryo <sup>2)</sup>	Total number <sup>2)</sup>	Ratio of ES origin (%)
C-6 <sup>3)</sup>	C-6-1	BALB/cA	BALB/cA	3	27	0	27	100
C-6 <sup>3)</sup>	C-6-2	BALB/cA	BALB/cA	7	16	37	53	30
C-6 <sup>3)</sup>	C-6-3	BALB/cA	BALB/cA	9	16	44	60	27
C-6 <sup>3)</sup>	C-6-4	BALB/cA	BALB/cA	6	6	36	42	14
C-6 <sup>3)</sup>	C-6-6	BALB/cA	BALB/cA	6	48	0	48	100
C-6 <sup>3)</sup>	CY-1	BALB/cA	BALB/cA	4	30	0	30	100
C-2 <sup>4)</sup>	B1	C57BL/6N	C57BL/6N	2	14	0	14	100
C-2 <sup>4)</sup>	B12	C57BL/6N	C57BL/6N	2	11	6	17	65
C-2 <sup>4)</sup>	B13	C57BL/6N	C57BL/6N	1	2	2	4	50
Cr-A-3 <sup>3)</sup>	CY-2	C3H/HeN	BALB/cA	5	4	41	45	9
Cr-A-3 <sup>3)</sup>	CY-3	C3H/HeN	BALB/cA	2	0	15	15	0
DB1 <sup>3)</sup>	DB1-1	C57BL/6J	DBA/1J	2	11 <sup>5)</sup>	0	11	100

1) Embryonic stem cells of the present invention: C-6 and C-2 were derived from C3H/HeN,

CrA/3 was derived from BALB/c

DB1 was derived from DBA/1J

- 2) Number of mice                      3) Aggregation method                      4) Injection method  
 5) Identified by PCR for CSA genotyping

The embryonic stem cell of the present invention are derived from genetically pure inbred strains, and therefore, extremely useful for close genetic researches. For example, they are useful for the production of knockout mice and analyses of cell genealogical tree, as well as research on the mechanism of embryonic stem cell establishment. In particular, as for the embryonic stem cells derived from DBA/1J and C3H strains and the like, a chimera mouse can be prepared using a non-treated embryonic stem cell, e.g., not introduced with a gene marker, and the behavior of the cells in the tissue of the chimera mouse, that are derived from the embryonic stem cells, can be analyzed at tissue level using

the monoclonal antibodies described in the U.S. Patent No. 5,552,287.

**Claims**

- 5 1. An embryonic stem cell derived from an inbred mouse strain.
2. An embryonic stem cell derived from a mouse strain selected from the group consisting of C3H/HeN, C57BL/6N, DBA/1J, and BALB/c.
- 10 3. Embryonic stem cell C-2 having the accession number FERM BP-5933 which is derived from mouse strain C3H/HeN.
4. Embryonic stem cell C-6 having the accession number FERM BP-5934 which is derived from mouse strain C3H/HeN.
- 15 5. Embryonic stem cell DB1 having the accession number FERM BP-5931 which is derived from mouse strain DBA/1J.
6. Embryonic stem cell B6-26 having the accession number FERM BP-5935 which is derived from mouse strain C57BL/6N.
- 20 7. Embryonic stem cell Cr/A-3 having the accession number FERM BP-5932 which is derived from mouse strain BALB/c.

25

30

35

40

45

50

55

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 808 896 A3**

(12)

**EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:  
22.04.1998 Bulletin 1998/17

(51) Int. Cl.<sup>6</sup>: C12N 5/00, C12N 5/06

(43) Date of publication A2:  
26.11.1997 Bulletin 1997/48

(21) Application number: 97108184.9

(22) Date of filing: 21.05.1997

(84) Designated Contracting States:  
DE FR GB

(30) Priority: 21.05.1996 JP 125533/96

(71) Applicant:  
The Institute of Physical and Chemical Research  
(RIKEN)  
Wako-shi, Saitama 351-01 (JP)

(72) Inventors:

- Kusakabe, Moriaki  
Koyadai, 3-1-1, Tsukuba-shi, Ibaragi 305 (JP)
- Kamon, Toshio  
20-14, Aobadai 2-chome, Meguro-ku Tokyo 153 (JP)

(74) Representative:

Godemeyer, Thomas, Dr.  
Hauptstrasse 58  
51491 Overath (DE)

**(54) Embryonic stem cells from mice**

(57) Embryonic stem cells derived from an inbred mouse strain selected from the group consisting of C3H/HeN, C57BL/6N, DBA/1J, and BALB/c strains, for example, C-2 cell having the accession number FERM BP-5933 and C-6 cell having the accession number FERM BP-5934 are disclosed. The embryonic stem cells of the present invention are derived from genetically complete inbred strains, and therefore, they are extremely useful for close genetic researches. For example, they are useful for the production of knockout mice as well as for analyses of cell genealogical tree and mechanism of embryonic stem cell establishment.

EP 0 808 896 A3

**BEST AVAILABLE COPY**



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 97 10 8184

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (In ICL6)
X	H. SUEMORI AND N. NAKATSUJI: "Establishment of the embryo-derived stem (ES) cell lines from mouse blastocysts: effects of the feeder cell layer" DEVELOPMENT. GROWTH AND DIFFERENTIATION, vol. 29, no. 2, 1987, NAGOYA, JP, pages 133-139, XP002056817	1	C12N5/00 C12N5/06
A	* the whole document *	2,6	
X	M.L. ROACH ET AL.: "A new embryonic stem cell line from DBA/11acJ mice allows genetic modification in a murine model of human inflammation" EXPERIMENTAL CELL RESEARCH, vol. 221, 1995, NEW YORK, NY, US, pages 520-525, XP002056818	1	
A	* the whole document *	2,5	
X	E. KAWASE ET AL.: "Strain difference in establishment of mouse embryonic stem (ES) cell lines" INTERNATIONAL JOURNAL OF DEVELOPMENTAL BIOLOGY, vol. 38, 1994, LEIOA, BILBAO, SP, pages 385-390, XP002056819	1	TECHNICAL FIELDS SEARCHED (In ICL6) C12N
A	* the whole document *	2,6,7	
P,X	N. NOBEN-TRAUTH ET AL.: "Efficient targeting of the Il-4 gene in a BALB/c embryonic stem cell line" TRANSGENIC RESEARCH, vol. 5, 1996, LONDON, GB, pages 487-491, XP002056820	1	
	* the whole document *		
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 25. February 1998	Examiner Mateo Rosell, A.M.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03/82 (P04C01)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 97 10 8184

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.8)
P,X	H. KITANI ET AL., : "Isolation of germline-transmissible embryonic stem (ES) cell line from C3H/He mice" ZOOLOGICAL SCIENCE, vol. 13, 1996, TOKYO, JP, pages 865-871, XP002056821 * the whole document *	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 25 February 1998	Examiner Mateo Rosell, A.M.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03/92 (P04C01)

**This Page Blank (uspto)**